BYRUMEN™

*PELLETED VITAMINS AND MINERALS WITH **MONECO**™ FOR DAIRY CATTLE, BEEF CATTLE, CALVES AND GOATS.*

Please note feed additives maybe added (conditions apply) including Flaveco, ProN8ure, Toxibind, Tyleco and some prescribed medications but only as directed by a veterinarian. See separate datasheets for details.


PRESENTATION: An attractive pellet for even distribution in feed.

ACTIVE CONSTITUENTS: Each 50 g contains:

		NRC #			NRC #
VITAMIN A	12,500 I.U.	17%	COBALT	10 mg	50%
VITAMIN D_3	2,500 I.U.	12%	COPPER	250 mg	100%
VITAMIN E	100 mg	20%	MANGANESE	250 mg	50%
IODINE	3 mg	25%	ZINC	1,500 mg	100%
SELENIUM	5 mg	70%	MONENSIN SODIUM as MONECO™	250 mg	

(These levels of active constituents are minimal levels).

 # Percentage of the National Research Council (NRC) 2001 requirements for a 680 kg cow producing greater than 35 L of milk per day.
NB - some constituents corrected for active bioavailability.



PROPERTIES: **BYRUMEN™** is a concentrate of vitamins and minerals in pellet form for dairy and beef cattle, calves and goats. Vitamins and minerals improve nutrient value, build health and optimise performance.
Moneco is added for improved feed efficiency in feedlot cattle, increased milk production under some conditions in dairy cows, as an aid in reducing the severity of non-clinical ketosis in lactating dairy cows, as an aid in the prevention of coccidiosis in cattle caused by *Eimeria zuernii* and *Eimeria bovis*, and as an aid in the prevention of coccidiosis in goats.

DOSAGE AND ADMINISTRATION: **BYRUMEN™** is best fed by mixing with other feed such as grain and meals. Add early in the feed mixing process to ensure thorough mixing.

Dairy and Beef Cattle: 50 g/head/day

Calves and Goats: 5 g/50 kg bodyweight

CAUTION: **Feeding BYRUMEN to lactating cows may result in decreased milk fat percentage but there is no effect on total fat yield. DO NOT USE in goats producing milk for human consumption. Horses, other equines or dogs should not be fed or allowed access to feed containing MONECO. Ingestion in these species may be fatal. The recommended levels of MONECO must not be exceeded. High levels may cause decreased feed intake, poor growth, diarrhoea and death.**

WITHHOLDING PERIODS: **CATTLE: MEAT AND MILK – NIL.**
GOATS:MEAT: NIL. MILK: DO NOT USE Byrumen in goats which are producing or may produce milk or milk products for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: S6

REGULATORY STATUS: EXEMPT.

PACK SIZE: 20 kg (contains 400 x 50 g doses or contains 4,000 x 5 g doses).

Australia-Wide Toll Free: 1800 801 201
Email: **info@iahp.com.au**
Website: **www.iahp.com.au**

® The Australia/World symbol is registered trademark of International Animal Health Products P/L
™ Byrumen and Moneco are trademarks of International Animal Health Products P/L.
© Copyright 1995 – 2026. All Rights Reserved.

10/2026