



## **DOTECO 1000™**

*AN AID IN THE PREVENTION OF COCCIDIOSIS OF BROILERS, REPLACEMENT PULLETS AND TURKEYS.*

**PRESENTATION:** Cream to light brown free-flowing powder free of particulate matter.

**ACTIVE CONSTITUENTS:** DINITOLMIDE (3,5-DINITRO-ORTHO-TOLUAMIDE) 1000 g/kg.

**PROPERTIES:** DOTECO™ 1000 is effective as an aid in the prevention of coccidiosis of broilers, replacement pullets and turkeys.

### **DOSE AND ADMINISTRATION:**

Use continuously for up to 14 days or as directed by a Veterinarian. Consult a Veterinarian for advice relating to coccidiosis control programs.

DOTECO 1000™ should be premixed with 5-10 kg of feed and then thoroughly mixed with the bulk feed.

**Broilers and Replacement Pullets:** Add DOTECO 1000™ at the rate of 125 grams/tonne of finished feed.

**Turkey Growers:** Add DOTECO 1000™ at the rate of 190 grams/tonne of finished feed.

### **WITHHOLDING PERIODS:**

**MEAT: NIL.**

**EGGS: DO NOT USE in laying hens (chickens/turkeys) or in replacement pullets over 14 weeks of age.**

**TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):** This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

**POISONS SCHEDULE:** NOT APPLICABLE

**REGULATORY STATUS:** AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

**PACK SIZES:** 25 kg